

<b>WARNING</b>	Any malpractice or any attempt to commit any kind of malpractice in the Examination will <b>DISQUALIFY THE CANDIDATE</b> .		
<b>PAPER – I PHYSICS &amp; CHEMISTRY-2019</b>			
Version Code	<b>A1</b>	Question Booklet Serial Number:	<b>5104300</b>
Time: 150 Minutes	Number of Questions : 120		Maximum Marks : 480
Name of the Candidate			
Roll Number			
Signature of the Candidate			
<b>INSTRUCTION TO CANDIDATES</b>			
<ol style="list-style-type: none"> <li>1. Please ensure that the <b>VERSION CODE</b> shown at the top of this Question Booklet is same as that shown in the <b>OMR Answer Sheet</b> issued to you. If you have received a Question Booklet with a different Version Code, please get it replaced with a Question Booklet with the same Version Code as that of OMR Answer Sheet from the Invigilator. <b>THIS IS VERY IMPORTANT.</b></li> <li>2. Please fill the items such as Name, Roll Number and Signature in the columns given above. Please also write Question Booklet Serial Number given at the top of this page against item 3 in the OMR Answer Sheet.</li> <li>3. This Question Booklet contains 120 questions. For each question five answers are suggested and given against (A), (B), (C), (D), and (E) of which only one will be the '<b>Most Appropriate Answer.</b>' Mark the bubble containing the letter corresponding to the '<b>Most Appropriate Answer</b>' in the OMR Answer Sheet, by using either <b>Blue or Black Ball Point Pen only.</b></li> <li>4. <b>Negative Marking:</b> In order to discourage wild guessing the score will be subjected to penalization formula based on the number of right answers actually marked and the number of wrong answer marked. Each correct answer will be awarded <b>FOUR</b> marks. <b>ONE</b> mark will be deducted for each incorrect answer. More than one answer marked against a question will be deemed as incorrect answer and will be negatively marked.</li> <li>5. Please read the instructions in the OMR Answer Sheet for marking the answers. Candidates are advised to strictly follow the instruction contained in the OMR Answer Sheet.</li> </ol>			
<b>IMMEDIATELY AFTER OPENING THE QUESTION BOOKLET, THE CANDIDATE SHOULD VERIFY WHETHER THE QUESTION BOOKLET CONTAINS ALL THE 120 QUESTIONS IN SERIAL ORDER. IF NOT, REQUEST FOR REPLACEMENT.</b>			
<b>DO NOT OPEN THE SEAL UNTIL THE INVIGILATOR ASKS YOU TO DO SO.</b>			

**SEAL**

WARNING: If the Examination will DISQUALIFY THE CANDIDATE.	
<b>TAPER - I PHYSICS &amp; CHEMISTRY -2019</b>	
Question Booklet Serial Number:	<b>A1</b>
Candidate's Booklet Serial Number:	<b>3104300</b>
Time: 150 Minutes	Number of Questions: 120
Name of the Candidate	Roll Number
Signature of the Candidate	Date

**INSTRUCTIONS TO CANDIDATES**

1. Please ensure that the PATTERN CODE shown at the top of the Question Booklet is same as that shown in the OMR Answer Sheet issued to you. If you find a discrepancy, please inform the invigilator immediately. Do not proceed with the question booklet until you are given the go-ahead signal by the invigilator. The question booklet will be given to you as per the PATTERN CODE shown at the top of the Question Booklet.

2. The Question Booklet contains 120 questions. For each question five answers are suggested and given against (A), (B), (C), (D), and (E) of which only one will be the 'Most Appropriate Answer'. Mark the bubble containing the correct response to the 'Most Appropriate Answer' in the OMR Answer Sheet by using either Blue or Black Ball Point Pen only.

3. Negative Marking: In order to discourage wild guessing the correct will be subjected to negative marking. For each question, one mark will be awarded for a correct answer and one mark will be deducted for an incorrect answer. If you are not sure of the correct answer, you may leave the question blank. No marks will be awarded for an unattempted question. The question will be deemed as incorrect answer and will be negatively marked.


4. Please refer to the instructions in the OMR Answer Sheet for marking the answers. Candidates are advised to strictly follow the instructions mentioned in the OMR Answer Sheet.

**BLANK PAGE**



PLEASE ENSURE THAT THIS QUESTION BOOKLET CONTAINS 120  
QUESTIONS SERIALLY NUMBERED FROM 1 TO 120. PRINTED PAGES 32.

- The dimensions for pressure is  
(A)  $MLT^{-2}$  (B)  $ML^{-1}T^{-2}$  (C)  $M^{-1}L^{-1}T^{-2}$   
(D)  $ML^{-1}T^{-1}$  (E)  $MLT$
- The magnitude of deceleration required for a body, moving at a speed of 10 m/s to come to a complete halt at a distance of 100 m is  
(A)  $20 \text{ m/s}^2$  (B)  $10 \text{ m/s}^2$  (C)  $2 \text{ m/s}^2$   
(D)  $0.5 \text{ m/s}^2$  (E)  $1 \text{ m/s}^2$
- An accurate measurement implies that  
(A) the spread of the readings are broad around the mean value  
(B) the spread of the readings are narrow around the mean value  
(C) the mean value of the readings is always lower than the true value  
(D) the mean value of the readings is always higher than the true value  
(E) the closeness of the mean of the readings to the true value
- The following plot gives the variation of acceleration ( $\text{m/s}^2$ ) with time (s) for an object that started from rest at time  $t = 0 \text{ s}$ . The velocity at time  $t = 15 \text{ s}$  ( $V_{15}$ ) and at 25 s ( $V_{25}$ ), respectively are



- (A)  $V_{15} = 50 \text{ m/s}$  and  $V_{25} = 0 \text{ m/s}$  (B)  $V_{15} = 100 \text{ m/s}$  and  $V_{25} = 150 \text{ m/s}$   
(C)  $V_{15} = 50 \text{ m/s}$  and  $V_{25} = 25 \text{ m/s}$  (D)  $V_{15} = 100 \text{ m/s}$  and  $V_{25} = 25 \text{ m/s}$   
(E)  $V_{15} = 75 \text{ m/s}$  and  $V_{25} = 50 \text{ m/s}$

Space for rough work

5. An object, moving with velocity 5 m/s, undergoes an acceleration of  $1 \text{ m/s}^2$  at time  $t = 0$ . If the object has a mass of 1 kg, the kinetic energy (KE) of the object at time  $t = 5 \text{ s}$  is
- (A) KE = 12.5 Joules      (B) KE = 20 Joules      (C) KE = 30 Joules  
 (D) KE = 50 Joules      (E) KE = 0 Joules
6. The variation of speed (in m/s) of an object with time (in seconds) is given by the expression  $V(t) = V_0 - 5t + 5t^2$
- (A) At time  $t = 0 \text{ s}$ , the instantaneous acceleration is zero  
 (B) At time  $t = 0 \text{ s}$ , there is a deceleration of the object  
 (C) At time  $t = 1 \text{ s}$ , the object is at rest  
 (D) At time  $t = 1 \text{ s}$ , the instantaneous acceleration is zero  
 (E) The distance travelled by the object at time  $t = 1 \text{ s}$  is  $V_0 \text{ m}$
7. A boat is moving from the east bank to the west bank on a south flowing river. If the speed of the boat is 4 km/h and that of the river is 3 km/h. If the width of the river is 2 km, the distance travelled by the boat is
- (A) 5 km      (B) 4 km      (C) 3 km  
 (D) 2.5 km      (E) 2 km
8. A bead is tied on one end of a stiff rope of length 1 m. With the other end of the rope as the center, the rope is rotated in such a way that the bead completes 10 revolutions per second. The centripetal acceleration of the bead is
- (A)  $400 \pi^2 \text{ m/s}^2$       (B)  $200 \pi^2 \text{ m/s}^2$       (C)  $400 \text{ m/s}^2$   
 (D)  $200 \text{ m/s}^2$       (E)  $100 \text{ m/s}^2$

---

Space for rough work



9. The electric field of an electromagnetic wave in free space is given by

$$\vec{E} = 5 \sin\left(\frac{2\pi}{3}z - \omega t\right) \hat{y} \text{ V/m. Which of the following statements is correct?}$$

- (A) The wave propagates along  $\hat{y}$
- (B) The wave vector is given  $\vec{k} = \frac{2\pi}{3} \hat{z}$
- (C) The wavelength of the electromagnetic wave is  $\frac{1}{3}$  m
- (D) The corresponding magnetic field is  $\vec{B} = \frac{5}{c} \cos\left(\frac{2\pi}{3}z - \omega t\right) \hat{x}$  T
- (E) The frequency of the wave is approximately  $10^6$  Hz
10. The radiation produced by a 100 W bulb has the following property
- (A) The radiation is in the form of an electromagnetic wave which carries energy but not momentum
- (B) The radiation is in the form of an electromagnetic wave which carries momentum but not energy
- (C) The radiation is in the form of an electromagnetic wave which carries both energy and momentum
- (D) The radiation neither carries energy nor momentum
- (E) The intensity of radiation is independent of the distance from source
11. A parallel plate capacitor (of capacitance  $C$ ) with circular plates of radius  $r_0$  located at positions  $\pm \alpha$ , is connected in series with a resistor  $R$  and is charged by a battery of voltage  $V$ . Consider a circular loop  $L$  of radius  $\frac{r_0}{2}$  parallel to the capacitor plates is located at the center. Which of the following statements is correct?
- (A) The charge on the capacitor at time  $t$  is  $q(t) = CR\left(1 - e^{-t/(CR)}\right)$
- (B) The charge on the capacitor at time  $t$  is  $q(t) = CV\left(1 - e^{-t/(CR)}\right)$
- (C) The flux through the loop  $L$  is independent of the area enclosed by it
- (D) The magnetic field is directed orthogonal to the loop  $L$
- (E) The magnetic field is directed along the loop  $L$

---

Space for rough work

12. A monochromatic light of frequency  $\nu = \frac{1}{6.63} \times 10^{16}$  Hz is produced by a laser. The power emitted is  $P = 10^{-2}$  W. The average number of photons per second emitted by the source is
- (A)  $\frac{1}{(6.63)^2} \times 10^{16}$       (B)  $(6.63)^2 \times 10^{20}$       (C)  $(6.63)^2 \times 10^{16}$   
 (D)  $10^{20}$       (E)  $10^{16}$
13. The work function of three photosensitive materials used to build photoelectric devices are given as: Sodium (2.75 eV), copper (4.65 eV) and gold (5.1 eV). Which of the following statements is **correct**. (The frequency of visible light lies in the range  $4 \times 10^{14}$  Hz to  $8 \times 10^{14}$  Hz)?
- (A) Devices built by copper and gold can operate with visible light  
 (B) Devices built using sodium can operate with ultraviolet light  
 (C) All the devices can operate with infrared light  
 (D) All the devices can operate with visible light  
 (E) No device can operate with visible light
14. An object is placed at 9 cm in front of a concave mirror of radius of curvature 12 cm. The following statement is **true**
- (A) The image is formed 36 cm behind the mirror  
 (B) The image is 36 cm in front of the mirror  
 (C) The image is magnified, virtual and erect  
 (D) The image is magnified, real and erect  
 (E) The image is magnified, real and inverted
15. An optician prescribes a lens of power +2.5 D. The focal length of the lens in water is (Refractive indices of the lens and water are respectively 1.5 and 1.33)
- (A) 40 cm      (B) 2660/17 cm      (C) 17/2660 cm  
 (D) 3000/17 cm      (E) 17/3000 cm

---

Space for rough work



16. In a single slit diffraction (of width  $\alpha$ ) by a monochromatic source of wavelength  $\lambda$  the first minimum of the intensity distribution occurs at an angle
- (A)  $\frac{\lambda}{\alpha}$                       (B)  $\frac{\lambda}{2\alpha}$                       (C)  $\frac{\alpha}{\lambda}$   
 (D)  $\frac{\alpha}{2\lambda}$                       (E)  $\frac{\pi}{4}$
17. A monochromatic source of wavelength 600 nm was used in Young's double slit experiment to produce interference pattern.  $I_1$  is the intensity of light at a point on the screen where the path difference is 150 nm. The intensity of light at a point where the path difference is 200 nm is given by
- (A)  $\frac{1}{2} I_1$                       (B)  $\frac{3}{2} I_1$                       (C)  $\frac{2}{3} I_1$   
 (D)  $\frac{3}{4} I_1$                       (E)  $\frac{4}{3} I_1$
18. The Brewster's angle for air to water interface is
- (A)  $\tan^{-1}(1.33)$                       (B)  $\sin^{-1}(1.33)$                       (C)  $\cos^{-1}(1.33)$   
 (D)  $\tan^{-1}\left(\frac{1}{1.33}\right)$                       (E)  $\sin^{-1}\left(\frac{1}{1.33}\right)$
19. A TV transmitting antenna is 81 m tall. It has a half-power beam width of 10 degrees. If the receiving antenna is at the ground level, the service area covered by the transmitter is determined by
- (A) the half-power beam width, the height of the transmitter and the radius of the earth  
 (B) the height of the transmitter and the radius of the earth  
 (C) the half-power beam width and the radius of the earth  
 (D) the height of the transmitter and the half-power beam width  
 (E) the height of the transmitter

---

Space for rough work



20. In the amplitude modulation mode of transmission, the normal speech signal is with a maximum frequency of 5 kHz. If the carrier frequency is 200 kHz, the modulated signal will have the frequencies varying between  
(A) 190 kHz to 210 kHz (B) 195 kHz to 205 kHz (C) 195 kHz to 200 kHz  
(D) 200 kHz to 205 kHz (E) 199.5 kHz to 200.5 kHz
21. For signal transmission, modulation is necessary  
(A) to reduce distortion of the signal  
(B) to modify the frequency content of the signal  
(C) to mask the signal information from enemy  
(D) to radiate the signal to a large distance using antennas  
(E) to make it easy to amplify the signal
22. A light emitting diode is  
(A) a n-p-n type semiconductor with a forward bias  
(B) a p-n-p semiconductor with a reverse bias  
(C) a p-n-p semiconductor with a forward bias  
(D) a p-n semiconductor with a reverse bias  
(E) a p-n semiconductor with a forward bias
23. In the context of p-n junction, select the **correct** statement from the following  
(A) The barrier potential remains constant under forward bias  
(B) The width of the depletion region depends on the doping level in the p-type and n-type regions  
(C) Under forward bias condition, the p-n junction behaves like a pure resistor irrespective of bias voltage  
(D) Under reverse bias condition, the p-n junction behaves like a pure resistor irrespective of bias voltage  
(E) The barrier potential decreases under reverse bias

---

Space for rough work

24. The radius of gyration about an axis through the center of a hollow sphere with external radius  $a$  and internal radius  $b$  is

(A)  $\sqrt{\frac{2(a^3 - b^3)}{5(a^5 - b^5)}}$

(B)  $\sqrt{\frac{1(a^4 - b^4)}{4(a^2 - b^2)}}$

(C)  $\sqrt{\frac{1(a^5 - b^5)}{2(a^3 - b^3)}}$

(D)  $\sqrt{\frac{2(a^5 - b^5)}{5(a^3 - b^3)}}$

(E)  $\sqrt{\frac{5(a^4 - b^4)}{2(a^2 - b^2)}}$

25. A ball of mass 1 kg and radius 0.5 m, starting from rest rolls down on a  $30^\circ$  inclined plane. The torque acting on the ball at the distance of the 7 m from the starting point is close to

(Take acceleration due to gravity as  $10 \text{ m/s}^2$ )

(A) 0.25 N-m

(B) 0.7 N-m

(C) 0.5 N-m

(D) 0.4 N-m

(E) 1.4 N-m

26. If the radius of the earth suddenly decreases by half of its present value. Then the time duration of one day will be

(A) 6 hours

(B) 8 hours

(C) 12 hours

(D) 24 hours

(E) 48 hours

27. A hollow sphere and a solid sphere, of equal mass and equal radii roll down without slipping on an inclined plane. If the torque experienced by the hollow sphere and solid sphere are  $\tau_H$  and  $\tau_S$  respectively, then

(A)  $\tau_H < \tau_S$

(B)  $\tau_H > \tau_S$

(C)  $\tau_H = \tau_S$

(D)  $\tau_H = 0$

(E)  $\tau_S = 0$

28. A brick of mass 2 kg slides down an incline of height 5 m and angle  $30^\circ$ . If the coefficient of friction of the incline is  $\frac{1}{2\sqrt{3}}$ , the velocity of the block at the

bottom of the incline is

(Assume the acceleration due to gravity is  $10 \text{ m/s}^2$ )

(A) 5 m/s

(B) 50 m/s

(C) 7 m/s

(D) 0

(E) 10 m/s

---

Space for rough work



29. A spherically uniform planet of mass  $8 \times 10^{24}$  kg and of radius  $6 \times 10^6$  m is orbiting around the Sun. The escape velocity for the planet is close to  
(Take  $G = 6 \times 10^{-11}$  N-m<sup>2</sup>/kg<sup>2</sup>)
- (A) 11.2 km/s                      (B) 16 km/s                      (C) 4 km/s  
(D) 12.6 km/s                      (E) 1.6 km/s
30. Suppose two planets A and B revolve around a Sun in the galaxy. The semi-major axis of A and B are 1 and 5 AU (astronomical unit) respectively. If the period of revolution of A is 1 year, the period of revolution of B is
- (A) 1 year                      (B) 5 years                      (C) 11 years  
(D) 25 years                      (E) 125 years
31. The half-life of  ${}_{43}\text{Tc}^{99}$  is 6 hours. If 12 mg of  ${}_{43}\text{Tc}^{99}$  is injected to a patient, after 24 hours how much Tc will be left in the patient's body?
- (A) 0 mg                      (B) 0.75 mg                      (C) 3 mg  
(D) 6 mg                      (E) 12 mg
32. For atoms, which of the following statement is **correct**?
- (A) Heavier nuclei have more neutrons than protons  
(B) Heavier nuclei have equal number of protons and neutrons  
(C) Lighter nuclei have more neutrons than protons  
(D) Lighter nuclei have less number of neutrons than protons  
(E) For heavier nuclei, atomic mass varies as square of atomic number
33. Which of the following statements is **not** a property of nuclear force?
- (A) it is an attractive force  
(B) it is independent of interacting nucleons  
(C) it is a long range force  
(D) it is non-central force  
(E) it is a short range force
34. Which of the following elements you need to remove to form an isotone family?
- (A)  ${}_{8}\text{O}^{16}$                       (B)  ${}_{7}\text{N}^{15}$                       (C)  ${}_{6}\text{C}^{14}$   
(D)  ${}_{13}\text{Al}^{27}$                       (E)  ${}_{9}\text{F}^{17}$

---

Space for rough work



35. A magnetic field of 1 T applied at an angle  $\pi/3$  to the vertical direction is decreased to zero at a steady rate in one second. The magnitude of induced emf in a horizontally placed circular loop of radius 5 cm is given by
- (A)  $1.25\sqrt{3}\pi$  mV      (B)  $12.5\sqrt{3}\pi$  V      (C)  $1.25\pi$  mV  
 (D)  $12.5\pi$  V      (E)  $25\pi$  V
36. The dimension of mutual inductance is (Denote dimension of current as A)
- (A)  $ML^2T^2A^{-2}$       (B)  $ML^2T^{-2}A^{-2}$       (C)  $ML^{-2}T^2A^{-2}$   
 (D)  $ML^2T^{-3}A^{-1}$       (E)  $ML^2T^{-3}A^{-3}$
37. A pure inductor of inductance 0.1 H is connected to an AC source (of rms voltage) 220 V and angular frequency of 300 Hz. The rms current is
- (A)  $\frac{3}{22}$  A      (B)  $\frac{22}{3}$  A      (C)  $\frac{11}{150}$  A  
 (D)  $\frac{150}{11}$  A      (E)  $\frac{11}{3\pi}$  A
38. In an LCR series circuit (of inductance L, capacitance C and resistance R), the impedance is minimum when the angular frequency of the source is given by
- (A)  $\sqrt{LC}$       (B)  $\frac{1}{\sqrt{LC}}$       (C)  $\sqrt{\frac{L}{C}}$   
 (D)  $\sqrt{\frac{C}{L}}$       (E)  $\sqrt{LCR}$

---

Space for rough work

39. A Carnot engine is operating between a hot body and cold body maintained at temperatures  $T_1$  and  $T_2$  respectively. Consider the following three cases
- Case-I: The temperature of the hot body is changed to  $T_1 + \Delta T$  and cold body is at  $T_2$
- Case-II: The temperature of the hot body is at  $T_1$  and cold body is changed to  $T_2 + \Delta T$
- Case-III: The temperature of the hot body is at  $T_1$  and cold body is changed to  $T_2 - \Delta T$
- (A) The efficiency of the Carnot cycle is highest for case-I  
(B) The efficiency of the Carnot cycle is highest for case-II  
(C) The efficiency of the Carnot cycle is highest for case-III  
(D) The efficiency of case-II is higher than case-III  
(E) The efficiency of the Carnot cycle is same for all three cases
40. Some smoke is trapped in a small glass container and is viewed through a microscope. A number of very small smoke particles are seen in continuous random motion as a result of their bombardment by air molecules. If the mass of the smoke particle is about  $10^{12}$  times higher than that of an air molecule the average speed of a smoke particle is
- (A)  $10^6$  times the average speed of an air molecule  
(B)  $10^{-12}$  times the average speed of an air molecule  
(C)  $10^{12}$  times the average speed of an air molecule  
(D)  $10^{-6}$  times the average speed of an air molecule  
(E)  $10^{-10}$  times the average speed of an air molecule
41. The standard of length is maintained by a 1 meter long bar made up of a material having coefficient of linear expansion  $\alpha = 0.00001 \text{ } ^\circ\text{C}^{-1}$ . If the length of the bar were to be preserved to an accuracy of 1 part per million, what would be maximum allowed temperature variation?
- (A)  $\pm 0.01 \text{ } ^\circ\text{C}$                       (B)  $\pm 0.1 \text{ } ^\circ\text{C}$                       (C)  $\pm 0.001 \text{ } ^\circ\text{C}$   
(D)  $\pm 0.0001 \text{ } ^\circ\text{C}$                       (E)  $\pm 1.0 \text{ } ^\circ\text{C}$

---

Space for rough work



42. Inside the engine of an automobile, the cylinder compresses the air from approximately standard temperature and pressure to one-twentieth of the original volume and a pressure of 50 atm. What is the temperature of the compressed air?  
(A) 500 K (B) 682 K (C) 550 K  
(D) 1000 K (E) 200 K
43. A spring of natural length  $l$  and spring constant 50 N/m is kept on a horizontal frictionless table with one end attached to a rigid support. First the spring was compressed by 10 cm and then released to hit a ball of mass 20 g kept at a distance  $l$  from the rigid support. If after hitting the ball, the natural length of the spring is restored, what is the speed with which the ball moved?  
(Ignore the air resistance)  
(A) 5 m/s (B) 7 m/s (C) 25 m/s  
(D) 50 m/s (E) 2500 m/s
44. In a water container, an aluminum piece of volume  $0.5 \text{ m}^3$  is lowered through an external force, until it is completely submerged. In another identical water container, a lead piece of same volume was similarly submerged using the same amount of external force. The mass density of lead is 4 times larger than the mass density of the aluminum. If  $F_A$  and  $F_L$  are the buoyancy forces acting on aluminum and lead respectively, then which of the following statements is correct?  
(A)  $F_A > 4 F_L$  (B)  $F_L > 4 F_A$  (C)  $F_A > 2 F_L$   
(D)  $F_L > 2 F_A$  (E)  $F_L = F_A$
45. A boy formed a bubble and a liquid drop from the same soapy water. The pressure difference between inside and outside of the soap bubble is measured to be  $100 \text{ N/m}^2$ . If the radius of the droplet is half of the radius of the bubble, then the pressure difference between the inside and outside of the droplet is  
(A) 0 (B)  $50 \text{ N/m}^2$  (C)  $100 \text{ N/m}^2$   
(D)  $200 \text{ N/m}^2$  (E)  $400 \text{ N/m}^2$

---

Space for rough work



46. A mass  $m$ , suspended vertically by a massless ideal spring with spring constant  $k$ , is at rest. The mass is displaced upward by a height  $h$ . When released, the kinetic energy of the mass will be proportional to  
(Neglecting air resistance)
- (A) only  $h$
  - (B) only  $h^2$
  - (C)  $m$
  - (D) a linear combination of terms involving  $h$  and  $h^2$
  - (E)  $k$
47. Instantaneous power delivered to a damped harmonic oscillator (natural frequency is  $\omega_0$ ) by an external periodic force (driving frequency  $\omega$ ) under steady state conditions is
- (A) positive always
  - (B) negative always
  - (C) positive and negative with power integrated over a period being zero
  - (D) positive and negative with power integrated over a period being positive
  - (E) positive and negative with power integrated over a period being negative
48. The Q factor for a damped oscillator is given by the
- (A) Ratio of energy stored per cycle to the initial energy
  - (B) Ratio of energy dissipated per cycle to the initial energy
  - (C) Ratio of energy stored per cycle to the energy dissipated per cycle
  - (D) Ratio of energy dissipated per cycle to the energy stored per cycle
  - (E) Ratio of the damping coefficient to the natural frequency
49. A ball of mass  $m$  is projected upward with a speed  $v_0$ . The speed at a height  $h$  is  
(Neglecting air resistance)
- (A) independent of angle and direction of projection
  - (B) independent of mass, angle and the direction of projection
  - (C) dependent on the direction of projection
  - (D) dependent on the shape, size and mass of the ball and angle of projection
  - (E) dependent on mass of the ball but independent of the angle and direction of projection

---

Space for rough work

50. An object having a velocity 5 m/s is accelerated at the rate  $2 \text{ m/s}^2$  for 6 s. Find the distance travelled during the period of acceleration


- (A) 60 m (B) 25 m (C) 36 m  
(D) 66 m (E) 45 m

51. A vehicle moving at 36 km/hr is to be stopped by applying brakes in the next 5 m. If the vehicle weighs 2000 kg, determine the average force that must be applied on it

- (A)  $10^4 \text{ N}$  (B)  $2 \times 10^4 \text{ N}$  (C)  $3 \times 10^4 \text{ N}$   
(D)  $5 \times 10^3 \text{ N}$  (E)  $10^3 \text{ N}$


52. A block of mass 20 kg is suspended through two spring balances with negligible mass as shown in figure. What will be the readings in the upper and lower balance respectively?

- (A) 0 kg, 20 kg  
(B) 10 kg, 20 kg  
(C) 20 kg, 10 kg  
(D) 10 kg, 10 kg  
(E) 20 kg, 20 kg



53. Two masses connected in series with two massless strings are hanging from a support as shown in the figure. Find the tension in the upper string

- (A)  $m_1 g$   
(B)  $(m_1 - m_2)g$   
(C)  $m_2 g$   
(D)  $(m_1 + m_2)g$   
(E)  $(m_1 \times m_2)g$



Space for rough work




54. An electron, placed in an electric field, experiences a force  $F$  of 1 N. What are the magnitude and direction of the electric field  $E$  at the point where the electron is located ( $e = 1.6 \times 10^{-19}$  C)?
- (A)  $\frac{1}{e}$  N/C,  $F$  and  $E$  are along the same direction  
 (B)  $\frac{1}{e}$  N/C,  $F$  and  $E$  are against each other  
 (C)  $\frac{1}{e}$  N/C,  $F$  and  $E$  are perpendicular  
 (D)  $e$  N/C,  $F$  and  $E$  are against each other  
 (E)  $e$  N/C,  $F$  and  $E$  are perpendicular
55. The distance between two charges  $q_1 = +2 \mu\text{C}$  and  $q_2 = +8 \mu\text{C}$  is 15 cm. Calculate the distance from the charge  $q_1$  to the points on the line segment joining the two charges where the electric field is zero
- (A) 1 cm                                      (B) 2 cm                                      (C) 3 cm  
 (D) 4 cm                                      (E) 5 cm

56. Four point charges (with equal magnitude of charge of 5 C; but with different signs) are placed at four corners of a square of side 10 m. Assuming that the square is centered at the origin and the configuration of the charges are as given in the figure, the potential and the magnitude of electric field at the origin, respectively are

$$\left[ \text{Note : } k = \frac{1}{4\pi\epsilon_0} \right]$$

- (A) 0 V and 0 V/m  
 (B) 0 V and  $\frac{\sqrt{2}}{5}k$  V/m  
 (C)  $\frac{\sqrt{2}}{5}k$  V and  $\frac{\sqrt{2}}{5}k$  V/m  
 (D) 0 V and 5 V/m  
 (E)  $\frac{\sqrt{2}}{5}k$  V and 0 V/m




---


Space for rough work

57. A point dipole with dipole moment,  $\vec{p} = p_0 \hat{k}$ , is kept at the origin. An external electric field given by,  $\vec{E} = E_0(2\hat{i} - 3\hat{j} + 4\hat{k})$ , is applied on it. Which one of the following statements is **true**?

- (A) The force on the dipole is zero while torque rotates the dipole on the  $xy$ -plane
- (B) The force on the dipole moves it along the direction of electric field
- (C) The interaction energy between the dipole and electric field is zero
- (D) The potential due to the dipole alone on the  $xy$ -plane with  $z = 0$  depends on the value of  $p_0$
- (E) The application of the electric field orients the dipole along the  $-\hat{k}$  direction


58. Find the total capacitance and total charge on the capacitors

- (A)  $1.5 \text{ nF}, 9 \text{ nC}$
- (B)  $3.0 \text{ nF}, 18 \text{ nC}$
- (C)  $1.5 \text{ nF}, 4.5 \text{ nC}$
- (D)  $3.0 \text{ nF}, 9 \text{ nC}$
- (E)  $3.0 \text{ nF}, 4.5 \text{ nC}$



59. A circuit is made using  $R_1, R_2, R_3, R_4$  and a battery as shown in the following figure. Find the equivalent resistance of the given circuit and the current passing through  $R_3$

- (A)  $3 \Omega, \frac{1}{3} \text{ A}$
- (B)  $\frac{1}{3} \Omega, 27 \text{ A}$
- (C)  $\frac{2}{3} \Omega, \frac{21}{2} \text{ A}$
- (D)  $\frac{1}{3} \Omega, \frac{21}{2} \text{ A}$
- (E)  $\frac{2}{3} \Omega, \frac{21}{3} \text{ A}$




Space for rough work



60. Find the voltage and current passing through the resistor  $R_2$  shown in the following circuit

- (A) 3 V, 3 mA  
 (B) 1 V, 1 mA  
 (C) 3 V, 1 mA  
 (D) 1 V, 2 mA  
 (E) 2 V, 1 mA



61. The resistor  $R_1 = 3 \Omega$  and  $R_2 = 1 \Omega$  are connected in parallel to a 20 V battery. Find the heat developed in the resistor  $R_1$  in one minute

- (A) 600 J (B) 800 J (C) 6000 J  
 (D) 8000 J (E) 7000 J

62. The velocity and acceleration of a particle performing simple harmonic motion have a steady phase relationship. The acceleration shows a phase lead over the velocity in radians of

- (A)  $+\pi$  (B) 0 (C)  $+\pi/2$   
 (D)  $-\pi/2$  (E)  $-\pi$

63. Consider a driven damped mechanical oscillator is in resonance. Which of the following statements is **true**?

- (A) Driving frequency is twice the natural frequency of the oscillator  
 (B) Power transfer from the driving source to system is minimum  
 (C) Driving frequency is the same as the natural frequency of the oscillator  
 (D) The force damping the oscillations are at a minimum value  
 (E) The driving force is in phase with the displacement

64. A body undergoing simple harmonic motion has a maximum acceleration of  $8\pi \text{ m/s}^2$  and maximum speed of 1.6 m/s. What is the time period  $T$ ?

- (A) 0.1 seconds (B) 0.2 seconds (C) 0.3 seconds  
 (D) 0.4 seconds (E) 0.5 seconds

---

Space for rough work


65. A bat emits an ultrasonic sound wave at 33.0 kHz and receives an echo 0.1 s later. What is the distance of the bat from the object-producing echo? (Speed of sound in air is 330 m/s)
- (A) 10.0 m                      (B) 20.0 m                      (C) 33.0 m  
(D) 66.0 m                      (E) 16.5 m
66. A wave along a string has the following equation  $y = 0.05 \sin (28t - 2.0x)$  m (where  $t$  is in seconds and  $x$  is in meters). What are the amplitude, frequency and wavelength of the wave?
- (A) amplitude = 0.05 m, frequency = 4.456 Hz and wavelength = 3.518 m  
(B) amplitude = 0.05 m, frequency = 28 Hz and wavelength = 2.0 m  
(C) amplitude = 5.0 m, frequency = 4.456 Hz and wavelength = 3.518 m  
(D) amplitude = 0.05 m, frequency = 2.0 Hz and wavelength = 28 m  
(E) amplitude = 0.05 m, frequency = 3.456 Hz and wavelength = 4.518 m
67. A train sounds its whistle as it approaches an observer standing at a point near the track. The observer measures a frequency of 216 Hz as the train approaches and a frequency of 184 Hz as the train leaves. What is the frequency of its whistle?
- (A) 210 Hz                      (B) 190 Hz                      (C) 205 Hz  
(D) 202 Hz                      (E) 200 Hz

---

Space for rough work



68. The following figure is the combination of logic gates. The inputs are A and B. The output is C. Which one of the following choices gives the correct matching?



(A)

A	B	C
0	0	1
0	1	1
1	0	1
1	1	1

(B)

A	B	C
0	0	1
0	1	0
1	0	1
1	1	0

(C)

A	B	C
0	0	0
0	1	1
1	0	0
1	1	1

(D)

A	B	C
0	0	1
0	1	1
1	0	0
1	1	0

(E)

A	B	C
0	0	0
0	1	0
1	0	0
1	1	0

69. A particle of mass  $m$  and charge  $q$  with an initial velocity  $\vec{v}$  is subjected to a uniform magnetic field  $\vec{B}$  along the vertical direction. The particle will
- (A) follow a circular path if  $\vec{v}$  is along the vertical direction
  - (B) make helical motion if  $\vec{v}$  is along the horizontal direction
  - (C) make helical motion if  $\vec{v}$  is neither parallel nor orthogonal to  $\vec{B}$
  - (D) always make circular motion
  - (E) always make helical motion

Space for rough work

70. Consider a circular loop of radius  $R$  on the  $xy$ -plane carrying a steady current anticlockwise. The magnetic field at the center of the loop is given by

- (A)  $\frac{\mu_0}{2R} I \hat{x}$                       (B)  $\frac{\mu_0}{2R} I \hat{y}$                       (C)  $\frac{\mu_0}{2R} I \hat{z}$   
(D)  $\frac{\mu_0}{R} I \hat{x}$                       (E)  $\frac{\mu_0}{R} I \hat{y}$

71. Consider two parallel current carrying conductors separated by a distance. Which one of the following statements is **true**?

- (A) Currents flowing in same direction will lead to repulsion  
(B) Currents flowing in opposite directions will lead to attraction  
(C) The conductors will always attract each other  
(D) The conductors will always repel each other  
(E) Currents flowing in same direction will lead to attraction and opposite directions will lead to repulsion

72. The energy gap is much more in silicon than in germanium because

- (A) It has less number of electrons  
(B) It has high atomic mass number  
(C) Its crystal has much stronger bonds called ionic bonds  
(D) Its valence electrons are more tightly bound to their parent nuclei  
(E) Its valence electrons are more loosely bound to their parent nuclei

---

Space for rough work



73.  $\text{NO}^+$  has bond order  
(A) 2 (B) 2.5 (C) 3  
(D) 3.5 (E) 4
74. Which hydride amongst the following has the least boiling point?  
(A)  $\text{NH}_3$  (B)  $\text{PH}_3$  (C)  $\text{AsH}_3$   
(D)  $\text{SbH}_3$  (E)  $\text{BiH}_3$
75. Which of the following sets has Lewis acid behaviour for all the components?  
(A)  $\text{BF}_3$  (B)  $\text{BF}_3, \text{SiF}_4, \text{PF}_5$  (C)  $\text{SiF}_4, \text{PF}_5$   
(D)  $\text{BF}_3, \text{PF}_5$  (E) (A) and (B)
76. The carbon atoms in calcium carbide are held by  
(A) Ionic bonds  
(B) Two sigma bonds  
(C) Two sigma and one coordinate bond  
(D) One sigma and two  $\pi$  bonds  
(E) One sigma and one  $\pi$  bond
77. According to the VSEPR theory, the shape of  $\text{ClO}_3^-$  would be  
(A) Linear (B) Triangular planar (C) Pyramidal  
(D) Square planar (E) Angular
78. Acetic acid in liquid ammonia behaves as  
(A) Weaker acid than that in water  
(B) Stronger acid than that in water  
(C) Base acid  
(D) Neutral acid  
(E) (C) and (D)
79. The compound(s) that does(do) not exist is(are)  
(A)  $\text{BiF}_5$  (B)  $\text{PF}_5$  (C)  $\text{AsF}_5$   
(D)  $\text{SbF}_5$  (E) All the compounds exist

---

Space for rough work

80. Rare gases are sparingly soluble in water because of  
(A) Hydrogen bonding  
(B) Dipole–dipole interaction  
(C) Induced dipole–induced dipole interaction  
(D) Dipole–induced dipole interaction  
(E) (A) and (D)
81. An example of a non-stoichiometric oxide when heated is  
(A) BeO (B) ZnO (C) MgO  
(D) CaO (E) Li<sub>2</sub>O
82. The donor atom in EDTA are  
(A) Two N and two O (B) Two N and four O (C) Four N and two O  
(D) Three N and three O (E) Two N and three O
83. Hard acids prefer to combine with  
(A) Soft bases (B) Soft acids (C) Hard acids  
(D) Hard bases (E) Salts
84. Among the following, which species represents a pseudohalide?  
(A) CN<sup>-</sup> (B) CaO (C) I<sub>2</sub>  
(D) K<sub>2</sub>HgI<sub>4</sub> (E) BiOCl
85. PCl<sub>3</sub> is stored in a well stoppered bottle since  
(A) It decomposes in the presence of moisture  
(B) It is decomposed by light  
(C) It is highly volatile  
(D) It reacts with air to form POCl<sub>3</sub>  
(E) (A) and (C)
86. An orange solid (A) on heating gives a green residue (B), a colourless gas (C) and water vapours. The dry gas (C) upon passing over heated Mg gave a white solid (D) which upon subsequent reaction with water gave a gas (E) that gave dense white fumes with HCl. Identify (D)  
(A) Fe(NH<sub>3</sub>)Cl<sub>2</sub> (B) CuN<sub>2</sub> (C) Mg<sub>3</sub>N<sub>2</sub>  
(D) NH<sub>4</sub>Cl (E) FeCl<sub>2</sub>

---

Space for rough work

87. On passing silent electric discharge through oxygen in an ozonizer, 5.5 mol % of oxygen is converted to ozone. How many moles of  $O_2$  and  $O_3$  result when 35 moles of  $O_2$  is originally present?
- (A) 33.0 (B) 34.4 (C) 35.0  
(D) 31.8 (E) 31.0
88. Carnallite is a mineral containing
- (A) K (B) Na (C) Mg  
(D) Fe (E) (A) and (C)
89. Maximum number of photons emitted by a bulb capable of producing monochromatic light of wavelength 550 nm is \_\_\_\_\_, if 100 V and 1 A is supplied for one hour.
- (A)  $1 \times 10^{24}$  (B)  $5 \times 10^{24}$  (C)  $1 \times 10^{23}$   
(D)  $5 \times 10^{23}$  (E)  $5 \times 10^{22}$
90. Which of the following is the correct unit of angular momentum of an electron in an orbital of an atom?
- (A) J s (B) J / s (C)  $W / s^2$   
(D) W s (E)  $J s^2$
91. Consider a *fcc* lattice made of a metal cation ( $M^{6+}$ ) and three oxide anions per unit cell. The resultant structure would have
- (A) 3D network of edge shared octahedra  
(B) 3D network of corner shared octahedra  
(C) 2D network of edge shared octahedra  
(D) 2D network of corner shared octahedra  
(E) 3D network of face shared octahedra

---

Space for rough work



92. The edge length of a solid possessing cubic unit cell is  $2\sqrt{2}r$  (structure I), based on hard sphere model, which upon subjected to a phase transition, a new cubic structure (structure II) having an edge length of  $\frac{4r}{\sqrt{3}}$  is obtained, where  $r$  is the radius of the hard sphere. Which of the following statements is **true**?
- (A) Density of the structure II is lower than structure I  
 (B) Density of structure II is higher than structure I  
 (C) The pore volume in structure I is 1.2 times higher than that of structure II  
 (D) The pore volume of both the structures are equal  
 (E) The octahedral voids in structure I is transformed into tetrahedral voids in structure II
93. An ideal gas "A" having volume of 1 L at 27 °C is kept in a container having movable piston and adiabatic walls in ambient condition. If 1.33 L atm of energy is supplied inside the system, find out the final temperature of the system?
- (A) 399 K (B) 499 K (C) 599 K  
 (D) 299 K (E) 450 K
94. A 5.2 L closed container contains some water and  $N_2(g)$  at 29 °C. The total pressure of the system and water tension are 1 atm and 0.04 atm, respectively. Upon electrolysis the liquid water inside completely, the final pressure of system was at 2 atm. What is number of moles of water that was present inside the container?
- (A)  $\frac{3.46}{RT}$  (B)  $\frac{5.2}{RT}$  (C)  $\frac{10.4}{RT}$   
 (D)  $\frac{0.208}{RT}$  (E)  $\frac{8.0}{RT}$

---

Space for rough work

95. A solution of methanol in water is 20 % by volume. If the solution and pure methanol have densities of  $0.964 \text{ kg L}^{-1}$  and  $0.793 \text{ kg L}^{-1}$ , respectively, find the per cent of methanol by weight?  
(A) 15.8 (B) 16.45 (C) 20  
(D) 14.8 (E) 17.6
96. The Henry's law constant for  $\text{O}_2$  dissolved in water is  $4.34 \times 10^4 \text{ atm}$  at certain temperature. If the partial pressure of  $\text{O}_2$  in a gas mixture that is in equilibrium with water is  $0.434 \text{ atm}$ , what is the mole fraction of  $\text{O}_2$  in the solution?  
(A)  $1 \times 10^{-5}$   
(B)  $1 \times 10^{-4}$   
(C)  $2 \times 10^{-5}$   
(D)  $1 \times 10^{-6}$   
(E)  $2 \times 10^{-6}$
97. The standard heat of formation of  $\text{CH}_4$ ,  $\text{CO}_2$  and  $\text{H}_2\text{O} (\text{l})$  are  $-76.2$ ,  $-394.8$  and  $-285.82 \text{ kJ mol}^{-1}$ , respectively. Heat of vaporization of water is  $44 \text{ kJ mol}^{-1}$ . Calculate the amount of heat evolved when  $22.4 \text{ L}$  of  $\text{CH}_4$ , kept under normal conditions, is oxidized into its gaseous products  
(A)  $802 \text{ kJ}$  (B)  $878.4 \text{ kJ}$  (C)  $702 \text{ kJ}$   
(D)  $788.4 \text{ kJ}$  (E)  $500 \text{ kJ}$
98. Acetic acid dimerizes when dissolved in benzene. As a result boiling point of the solution rises by  $0.36^\circ\text{C}$ , when  $100 \text{ g}$  of benzene is mixed with "X" g of acetic acid. In this solution, if experimentally measured molecular weight of acetic acid is  $117.8$  and molar elevation constant of benzene is  $2.57 \text{ K kg mol}^{-1}$ , what is the weight % and degree of dissociation (in %) of acetic acid in benzene?  
(A)  $1.62$  and  $98.3$  (B)  $0.81$  and  $98.3$  (C)  $0.5$  and  $86$   
(D)  $1$  and  $98.3$  (E)  $1.4$  and  $99$
99. At a certain temperature,  $2$  moles of  $\text{CO}$  and  $4$  moles of  $\text{Cl}_2$  gases were reacted to form  $\text{COCl}_2$  in a  $10 \text{ L}$  vessel. At equilibrium if one mole of  $\text{CO}$  is present then equilibrium constant for the reaction is  
(A)  $4$  (B)  $3.3$  (C)  $1$   
(D)  $2.5$  (E)  $4.5$

---

Space for rough work

100. The equilibrium constant for the reaction,  $N_2(g) + 3H_2 \rightleftharpoons 2NH_3(g)$  and  $2N_2(g) + 6H_2 \rightleftharpoons 4NH_3(g)$  are  $K_1$  and  $K_2$ , respectively. The relationship between  $K_1$  and  $K_2$  is

- (A)  $K_2 = K_1^2$                       (B)  $K_2 = K_1^{-2}$                       (C)  $K_1 = K_2^2$   
 (D)  $K_2 = \sqrt{K_1}$                       (E)  $K_1 = \sqrt{K_2}$


101. For a first order reaction,  $A(g) \rightarrow B(g)$  at  $35^\circ C$ , the volume of "A" left in the reaction vessel at various times are given below. [Given data:  $\log(5/4) = 0.0969$ ]

$t / \text{minutes}$	0	10	20	30	40
$V / \text{mL}$	25	20	15.7	12.5	9.6

What is the value of rate constant?

- (A)  $0.02231 \text{ min}^{-1}$                       (B)  $0.04231 \text{ min}^{-1}$                       (C)  $0.06231 \text{ min}^{-1}$   
 (D)  $0.08231 \text{ min}^{-1}$                       (E)  $0.1231 \text{ min}^{-1}$

102.  $E_{\text{cell}}$  of the following cell is



- (A)  $\frac{-2.303RT}{F}$                       (B)  $\frac{2.303RT}{F}$                       (C)  $\frac{-2.303RT}{2F}$   
 (D)  $\frac{2.303RT}{2F}$                       (E)  $\frac{RT}{2F}$

103. In a lead-acid battery, if 1 A current is passed to charge the battery for 1 h, what is the amount of  $PbSO_4$  converted to  $PbO_2$ ? (Given data:  $1 F = 96500 \text{ C mol}^{-1}$ )

- (A) 0.0373 moles                      (B) 0.0186 moles                      (C) 0.0093 moles  
 (D) 0.0268 moles                      (E) 0.0400 moles

Space for rough work



104. A fuel cell operates at constant current, with  $H_2$  fuel (1 bar) and  $O_2$  oxidant (1 bar). The electrolyte used is 0.001 M HCl and the product(s) of the reaction are confined inside the fuel cell. Which of the following is **true** about the electrolyte?
- (A) Boiling point of the electrolyte decreases with increase in the duration of fuel cell operation
  - (B) Boiling point of the electrolyte increases with increase in the duration of fuel cell operation
  - (C) Open circuit voltage of the fuel cell remains constant with increase in duration of operation
  - (D) Open circuit voltage of the fuel cell increases with increase in duration of operation
  - (E) Both (A) and (C)
105. The correct IUPAC name for methylisopropylacetylene is
- (A) 2-methyl-4-pentyne
  - (B) 4-methyl-2-pentyne
  - (C) isopropylmethylacetylene
  - (D) 3-methyl-4-pentyne
  - (E) 2-methyl-3-pentyne
106. Cyclohexylamine and aniline can be distinguished by
- (A) Hinsberg's test
  - (B) Carbylamine test
  - (C) Bromine test
  - (D) Beilstein's test
  - (E) Lassaigne's test
107. The compounds pyridine and planar cyclooctatetraene are \_\_\_\_\_, respectively
- (A) aromatic and non-aromatic
  - (B) aromatic and anti-aromatic
  - (C) aromatic and aromatic
  - (D) anti-aromatic and non-aromatic
  - (E) anti-aromatic and anti-aromatic

---


Space for rough work

108. Propylene on treatment with  $\text{HBr}/\text{H}_2\text{O}_2$  provides  
(A) 1-bromopropane (B) 2-bromopropane (C) 1,2-dibromopropane  
(D) 1-bromopropan-2-ol (E) 2-bromopropan-1-ol
109. \_\_\_\_\_ is a potent vasodilator.  
(A) Histamine (B) Serotonin (C) Codeine  
(D) Cimetidine (E) Aspirin
110. An invert sugar is  
(A) Isorotatory (B) Levorotatory (C) Dextrorotatory  
(D) Optically inactive (E) Mutarotatory
111. The strongest base among the following is  
(A)  $\text{NH}_2^-$  (B)  $\text{OH}^-$  (C)  $\text{CH}=\text{C}^-$   
(D)  $\text{CH}_3\text{CH}_2^-$  (E)  $\text{OEt}^-$
112. The neopentyl halide in ethanol yields alkenes by  $\text{E1}$  mechanism due to  
(A) low concentration of solvent  
(B) absence of base  
(C) it is a primary halide  
(D) steric factor which prevents  $\text{E2}$  mechanism  
(E) solvation effect
113. Arylbromides are not good candidates for  
(A) Wurtz-Fittig reaction  
(B) Fittig reaction  
(C) Friedel-Crafts reaction  
(D) Grignard reaction  
(E) Gabriel-phthalimide synthesis
114. Sulfonation of benzene with excess sulfuric acid provides  
(A) benzenesulfonic acid  
(B) *p*-benzenedisulfonic acid  
(C) *o*-benzenedisulfonic acid  
(D) *m*-benzenedisulfonic acid  
(E) decomposition of benzene

---

Space for rough work

115. The following reaction is named as



- (A) Reimer-Tiemann
- (B) Kolbe-Schmitt
- (C) Cannizzaro
- (D) Gattermann
- (E) Aldol

116. When  $C_6H_5COCOC_6H_5$  is reduced with  $LiAlH_4$ , the product formed has \_\_\_\_\_ stereoisomers.

- (A) 2
- (B) 3
- (C) 4
- (D) 6
- (E) 8

117. The compound which does not lead to benzoic acid by oxidation with  $KMnO_4$  is

- (A) toluene
- (B) benzyl alcohol
- (C) *n*-butylbenzene
- (D) *t*-butylbenzene
- (E) styrene

118. In the Hofmann rearrangement of primary amides having optically active group with S-configuration, the product amine has

- (A) R-configuration
- (B) S-configuration
- (C) Racemic mixture
- (D) Meso form
- (E) Achiral nature


Space for rough work



119. Benzonitrile can be prepared from benzaldehyde on treatment with

- (A)  $\text{NH}_3$
- (B)  $\text{NH}_3$  followed by hydrogenation with Ni
- (C)  $\text{NH}_2\text{OH}$
- (D)  $\text{NH}_2\text{OH}$  followed by dehydration with acetic anhydride
- (E) Hydrogen cyanide

120. The product formed in the below reaction is



Space for rough work

SEAL

BLANK PAGE

**PAPER – I PHYSICS & CHEMISTRY-2019 VERSION-A1**

1	B	31	B	61	D	91	B
2	D	32	A	62	C	92	A
3	E	33	C	63	C	93	A
4	B	34	D	64	D	94	A
5	D	35	C	65	E	95	B
6	B	36	B	66	A	96	A
7	D	37	B	67	E	97	A
8	A	38	B	68	E	98	A
9	B	39	C	69	C	99	B
10	C	40	D	70	C	100	A
11	E	41	B	71	E	101	A
12	E	42	B	72	D	102	A
13	B	43	A	73	C	103	B
14	E	44	E	74	B	104	A
15	B	45	C	75	A	105	B
16	A	46	D	76	D	106	C
17	A	47	D	77	C	107	B
18	A	48	C	78	B	108	A
19	A	49	B	79	A	109	A
20	B	50	D	80	D	110	B
21	D	51	B	81	A	111	D
22	E	52	E	82	B	112	D
23	B	53	D	83	D	113	E
24	D	54	B	84	A	114	D
25	B	55	C	85	A	115	C
26	A	56	B	86	D	116	B
27	B	57	A	87	B	117	D
28	C	58	A	88	E	118	B
29	D	59	B	89	A	119	D
30	C	60	E	90	A	120	B

*Handwritten signature*  
3/5



**WARNING**

Any malpractice or any attempt to commit any kind of malpractice in the Examination will **DISQUALIFY THE CANDIDATE**.

**PAPER – II MATHEMATICS-2019**

Version Code

**B1**Question Booklet  
Serial Number :**6127980**

Time: 150 Minutes

Number of Questions : 120

Maximum Marks : 480

Name of the Candidate

Roll Number

Signature of the Candidate

**INSTRUCTIONS TO CANDIDATES**

1. Please ensure that the **VERSION CODE** shown at the top of this Question Booklet is same as that shown in the **OMR Answer Sheet** issued to you. If you have received a Question Booklet with a different Version Code, please get it replaced with a Question Booklet with the same Version Code as that of OMR Answer Sheet from the Invigilator. **THIS IS VERY IMPORTANT.**
2. Please fill the items such as Name, Roll Number and Signature in the columns given above. Please also write Question Booklet Serial Number given at the top of this page against item 3 in the OMR Answer Sheet.
3. This Question Booklet contains 120 questions. For each question five answers are suggested and given against (A), (B), (C), (D), and (E) of which only one will be the '**Most Appropriate Answer.**' Mark the bubble containing the letter corresponding to the '**Most Appropriate Answer**' in the OMR Answer Sheet, by using either **Blue or Black Ball Point Pen** only.
4. **Negative Marking:** In order to discourage wild guessing the score will be subjected to penalization formula based on the number of right answers actually marked and the number of wrong answer marked. Each correct answer will be awarded **FOUR** marks. **ONE** mark will be deducted for each incorrect answer. More than one answer marked against a question will be deemed as incorrect answer and will be negatively marked.
5. Please read the instructions in the OMR Answer Sheet for marking the answers. Candidates are advised to strictly follow the instruction contained in the OMR Answer Sheet.

**IMMEDIATELY AFTER OPENING THE QUESTION BOOKLET, THE CANDIDATE SHOULD VERIFY WHETHER THE QUESTION BOOKLET CONTAINS ALL THE 120 QUESTIONS IN SERIAL ORDER. IF NOT, REQUEST FOR REPLACEMENT.**

**DO NOT OPEN THE SEAL UNTIL THE INVIGILATOR ASKS YOU TO DO SO.**

**SEAL**

<p>For information only: This is a blank page. The examination will be held on the date mentioned in the advertisement. The examination will be held on the date mentioned in the advertisement.</p>	
<p><b>PAPER - II - MATHEMATICS - 2019</b></p>	
<p>Question Booklet Serial Number</p>	<p>81</p>
<p>Time: 120 minutes</p>	<p>Number of Questions: 40</p>
<p>Name of the Candidate</p>	
<p>Roll Number</p>	
<p>Signature of the Candidate</p>	
<p><b>BLANK PAGE</b></p>	

**PLEASE ENSURE THAT THIS QUESTION BOOKLET CONTAINS 120  
QUESTIONS SERIALLY NUMBERED FROM 1 TO 120. PRINTED PAGES 32.**

1. The axis of the parabola  $x^2 + 6x + 4y + 5 = 0$  is  
(A)  $x = 0$  (B)  $y = 1$  (C)  $x + 3 = 0$   
(D)  $y = 4$  (E)  $y + 2 = 0$
2. The distance between the foci of the ellipse  $\frac{(x+2)^2}{9} + \frac{(y-1)^2}{4} = 1$  is  
(A)  $\sqrt{5}$  (B)  $2\sqrt{5}$  (C)  $3\sqrt{5}$   
(D)  $9\sqrt{5}$  (E)  $7\sqrt{5}$
3. The value of  $k$ , if the circles  $2x^2 + 2y^2 - 4x + 6y = 3$  and  $x^2 + y^2 + kx + y = 0$  cut orthogonally is  
(A) 2 (B) 3 (C) 4  
(D) 5 (E) 1
4. The circle passing through  $(1, -2)$  and touching the  $x$ -axis at  $(3, 0)$  also passes through the point  
(A)  $(2, -5)$  (B)  $(-5, -2)$  (C)  $(-2, 5)$   
(D)  $(-5, 2)$  (E)  $(5, -2)$
5. If  $\alpha$  and  $\beta$  are the roots of the equation  $x^2 + \alpha x + \beta = 0$ , then  
(A)  $\alpha = -1, \beta = -2$  (B)  $\alpha = 0, \beta = 1$  (C)  $\alpha = -2, \beta = 0$   
(D)  $\alpha = -2, \beta = 1$  (E)  $\alpha = 1, \beta = -2$

---

Space for rough work



6. If  $\vec{a} = (1, 1, -1)$ ,  $\vec{b} = (-1, 2, 1)$  and  $\vec{c} = (-1, 2, -1)$ , then  $|(\vec{a} + \vec{b}) \times (\vec{b} + \vec{c})|$  is  
 (A) 2 (B) 4 (C) 6  
 (D) 8 (E) 10
7. A particle is displaced from the point  $(2, 1, -1)$  to the point  $(4, 3, -4)$  by the force  $2i + 4j - 5k$ . Then the work done by the force is  
 (A) 16 (B) 27 (C) 36  
 (D) 48 (E) 52
8. The value of  $m$  if the vectors  $4i - 3j + 5k$  and  $mi - 4j + k$  are perpendicular, is  
 (A)  $\frac{-15}{4}$  (B)  $\frac{-17}{4}$  (C)  $\frac{-19}{4}$   
 (D) 0 (E)  $\frac{11}{4}$
9. If A and B are two matrices such that  $3A + B = \begin{pmatrix} 9 & 11 & 3 \\ 12 & 14 & 19 \end{pmatrix}$   
 and  $2A - 3B = \begin{pmatrix} -16 & 11 & 2 \\ -3 & -22 & 9 \end{pmatrix}$ . Then the matrix B is  
 (A)  $\begin{pmatrix} 6 & -1 & 0 \\ 3 & 8 & 1 \end{pmatrix}$  (B)  $\begin{pmatrix} 3 & -1 & 0 \\ 2 & 1 & 1 \end{pmatrix}$  (C)  $\begin{pmatrix} 8 & 0 & -1 \\ 3 & 1 & 2 \end{pmatrix}$   
 (D)  $\begin{pmatrix} 5 & 3 & -1 \\ 0 & 1 & 2 \end{pmatrix}$  (E)  $\begin{pmatrix} 1 & -3 & 4 \\ 3 & 0 & 2 \end{pmatrix}$

---

Space for rough work

10. If  $a, b$  and  $c$  are distinct reals and the determinant  $\begin{vmatrix} a^3+1 & a^2 & a \\ b^3+1 & b^2 & b \\ c^3+1 & c^2 & c \end{vmatrix} = 0$ , then the

product  $abc$  is

- (A)  $-1$  (B)  $0$  (C)  $1$   
 (D)  $2$  (E)  $3$

11. If  $(x, y, z)$  is the solution of the equations

$$x - y - 2z = 3$$

$$2x + y + 4z = 5$$

$$4x - y - 2z = 11$$

then the value of  $y$  equals

- (A)  $0$  (B)  $-1/2$  (C)  $-1/3$   
 (D)  $-1/4$  (E)  $-1$

12. If  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  is the inverse of the matrix  $\begin{pmatrix} 1 & 5 \\ 7 & -3 \end{pmatrix}$ , then  $d$  equals

- (A)  $-1/38$  (B)  $-7/38$  (C)  $3/38$   
 (D)  $5/38$  (E)  $9/38$

13. If  $f: \mathbb{R} \rightarrow \mathbb{R}$  is a function defined by  $f(x) = \sin x$ , then which of the following is true?

- (A)  $f$  is 1-1 but not onto  
 (B)  $f$  is onto but not 1-1  
 (C)  $f$  is both 1-1 and onto  
 (D)  $f$  is neither 1-1 nor onto  
 (E)  $f$  has finite number of zeros

---

Space for rough work

14. Consider the set  $M = \{1, 2, 3\}$  along with the relation  $R = \{(1, 2), (1, 1), (3, 1), (3, 4), (3, 3), (4, 3)\}$ . Which of the following statements is **true**?
- (A) The relation is symmetric but not transitive  
 (B) The relation is transitive but not symmetric  
 (C) The relation is both symmetric and transitive  
 (D) The relation is neither symmetric nor transitive  
 (E) The relation is reflexive

15. Let  $z_1 = 1 + i\sqrt{3}$  and  $z_2 = 1 + i$ , then  $\arg\left(\frac{z_1}{z_2}\right)$  is

- (A)  $\frac{5\pi}{12}$                       (B)  $\frac{7\pi}{12}$                       (C)  $\frac{11\pi}{12}$   
 (D)  $\frac{3\pi}{12}$                       (E) Not defined

16. The complex number  $\sqrt{2}\left[\sin\frac{\pi}{8} + i\cos\frac{\pi}{8}\right]^6$  represents

- (A)  $-i$                       (B)  $i$                       (C)  $1 - i$   
 (D)  $1 + i$                       (E)  $1 + 2i$

17. If  $z^2 + z + 1 = 0$ , where  $z$  is a complex number, then the value of

$$\left(z + \frac{1}{z}\right)^2 + \left(z^2 + \frac{1}{z^2}\right)^2 + \left(z^3 + \frac{1}{z^3}\right)^2 + \dots + \left(z^6 + \frac{1}{z^6}\right)^2 \text{ is}$$

- (A) 18                      (B) 54                      (C) 6  
 (D) 19                      (E) 12

---

Space for rough work



18. The value of  $\tan \left[ \sin^{-1} \frac{-1}{\sqrt{2}} \right]$  is
- (A) -1 (B) 0 (C) 1  
(D) Infinity (E) 2
19. If  $\sin^{-1} x + \cos^{-1} 2x = \frac{\pi}{6}$ , then the value of  $x$  is
- (A)  $1/2$  (B)  $\sqrt{3}/2$  (C)  $\sqrt{3}$   
(D) 1 (E)  $\sqrt{2}$
20. If  $x = 2 \cos t - \cos 2t$  and  $y = 2 \sin t - \sin 2t$ , then  $\frac{dy}{dx}$  at  $t = \frac{\pi}{2}$  is
- (A) -1 (B) 0 (C)  $1/2$   
(D) 1 (E) 3
21. The equation of the tangent to the curve given by  $x^2 + 2x - 3y + 3 = 0$  at the point (1, 2) is
- (A)  $4x - 3y - 2 = 0$  (B)  $3y - 4x - 2 = 0$  (C)  $4x + 3y + 2 = 0$   
(D)  $4x + 3y - 2 = 0$  (E)  $4y - 3x + 2 = 0$
22. The value of  $\lim_{x \rightarrow \infty} \frac{x^3 \sin\left(\frac{1}{x}\right) - 2x^2}{1 + 3x^2}$  is
- (A) 0 (B)  $\frac{1}{3}$  (C) -1  
(D)  $-\frac{2}{3}$  (E)  $-\frac{1}{3}$

---

Space for rough work

23. The maximum value of  $y = \left(\frac{1}{x}\right)^x$ ,  $x > 0$  is
- (A)  $e^{1/e}$  (B)  $e^e$  (C) 1  
 (D) Infinity (E) 0
24. The value of the integral  $\int_0^\pi \frac{\cos x}{1 + \sin^2 x} dx$  is
- (A) 0 (B) 1 (C)  $\frac{\pi}{2}$   
 (D)  $\pi$  (E)  $2\pi$
25. The area enclosed between the curves  $y = 2x^2 + 1$  and  $y = x^2 + 5$  is
- (A)  $4/3$  (B)  $8/3$  (C)  $16/3$   
 (D)  $32/3$  (E)  $1/3$
26. The solution of the differential equation  $5y dx = 2x dy$  passing through the point (1, 1) is
- (A)  $2 \ln x = 5 \ln y$  (B)  $5 \ln x = 2 \ln y$  (C)  $\ln(y + x) = 2$   
 (D)  $\ln(1 + xy) = 0$  (E)  $3 \ln x = 5 \ln y$
27. The area of the region bounded by the curves  $y = |x - 2|$ ,  $x = 1$ ,  $x = 3$  and  $y = 0$  is
- (A) 4 (B) 12 (C) 3  
 (D) 14 (E) 1

---

Space for rough work

28. If in a frequency distribution, the mean and median are 21 and 22 respectively, then its mode is  
(A) 22.0 (B) 20.5 (C) 25.5  
(D) 23.2 (E) 24.0
29. For two data sets, each of size 5, the variances are given to be 4 and 5 and the corresponding means are given to be 2 and 4, respectively. The variance of the combined data set is  
(A)  $\frac{15}{2}$  (B) 6 (C)  $\frac{13}{2}$   
(D)  $\frac{5}{2}$  (E)  $\frac{11}{2}$
30. If the mean of the first  $n$  odd numbers is  $\frac{n^2}{81}$ , then  $n$  equals  
(A) 9 (B) 18 (C) 27  
(D) 81 (E) 52
31. A bag contains 5 red balls and some blue balls. If the probability of drawing a blue ball is double that of red ball, the number of blue balls must be  
(A) 10 (B) 15 (C) 20  
(D) 25 (E) 30
32. A pair of fair dice are rolled together. The probability of getting a total of 8 is  
(A)  $\frac{1}{9}$  (B)  $\frac{5}{36}$  (C)  $\frac{7}{36}$   
(D)  $\frac{11}{36}$  (E)  $\frac{1}{36}$

---

Space for rough work



33. In a chess tournament, assume that your probability of winning a game is 0.3 against level 1 players, 0.4 against level 2 players and 0.5 against level 3 players. It is further assumed that among the players 50 % are at level 1, 25 % are at level 2 and the remaining are at level 3. Suppose that you win the game. Then the probability that you had played with level 1 player is
- (A) 0.3                      (B) 0.4                      (C) 0.5  
(D) 0.6                      (E) 0.2
34. A sum of Rs. 280 is to be used to award four prizes. If each prize after the first prize is Rs. 20 less than its preceding prize, then the value of the fourth prize is
- (A) 20                      (B) 40                      (C) 60  
(D) 80                      (E) 10
35. The coefficient of  $x^3$  in the expansion of  $(1 + x + 2x^2)(1 - 2x)^5$  is
- (A) -20                      (B) -40                      (C) -60  
(D) -80                      (E) -100
36. The constant term in the expansion of  $\left(x^2 - \frac{2}{x}\right)^6$  is
- (A) 60                      (B) 180                      (C) 240  
(D) 360                      (E) 420

---

Space for rough work

37. If the equation of the sphere through the circle  $x^2 + y^2 + z^2 = 9$ ;  $2x + 3y + 4z = 5$  and through the point  $(1, 2, 3)$  is  $3(x^2 + y^2 + z^2) - 2x - 3y - 4z = C$ , then the value of  $C$  is  
 (A) 11 (B) 22 (C) 36  
 (D) 41 (E) 54
38. The equation of the plane containing the line  $\frac{x-\alpha}{l} = \frac{y-\beta}{m} = \frac{z-\gamma}{n}$  is  $a(x-\alpha) + b(y-\beta) + c(z-\gamma) = 0$ , where  $al + bm + cn$  is equal to  
 (A) 1 (B) -1 (C) 2  
 (D) 8 (E) 0
39. Let  $f(x)$  and  $g(x)$  be two differentiable functions for  $0 \leq x \leq 1$  such that  $f(0) = 2$ ,  $g(0) = 0$ ,  $f(1) = 6$ . If there exists a real number  $c$  in  $(0,1)$  such that  $f'(c) = 2g'(c)$ , then  $g(1)$  is equal to  
 (A) 0 (B) -1 (C) 4  
 (D) -2 (E) 2
40. The equation of the tangent to the curve  $y = x + \frac{4}{x^2}$  that is parallel to the  $x$ -axis is  
 (A)  $y = 1$  (B)  $y = 2$  (C)  $y = 8$   
 (D)  $y = 0$  (E)  $y = 3$

---

Space for rough work

41. The number 81 is the coefficient of  $x^k$  in the binomial expansion of  $\left(x^2 + \frac{3}{x}\right)^4$ ,  $x \neq 0$ . Then the value of  $k$  equals
- (A) -2 (B) 2 (C) -4  
(D) 4 (E) 5
42. The possible number of arrangements starting with K of the word KALINGA is
- (A) 300 (B) 330 (C) 360  
(D) 390 (E) 370
43. A bag contains 3 black and 2 white balls. A ball is drawn at random and is put back in the bag along with one ball of the same colour. A ball is again drawn at random. What is the probability that it is white?
- (A)  $\frac{1}{5}$  (B)  $\frac{2}{5}$  (C)  $\frac{1}{6}$   
(D)  $\frac{1}{12}$  (E)  $\frac{2}{13}$
44. If A and B are two events associated with an experiment such that  $P(A \cup B) = P(A \cap B)$ , and  $P(A) = \frac{1}{3}$ , then  $P(B)$  equals
- (A) 0 (B)  $\frac{1}{3}$  (C)  $\frac{2}{3}$   
(D)  $\frac{1}{2}$  (E)  $\frac{2}{5}$
45. Three identical fair dice are rolled. The probability that the same number appears on each of them is
- (A)  $\frac{1}{3}$  (B)  $\frac{1}{6}$  (C)  $\frac{1}{36}$   
(D)  $\frac{1}{216}$  (E)  $\frac{1}{9}$

---

Space for rough work



46. Let  $\omega \neq 1$  be a cube root of unity and  $(1 + \omega)^7 = a + \omega$ . Then the value of  $a$  is  
 (A)  $\omega^2$  (B)  $\omega$  (C)  $1/2$   
 (D)  $1$  (E)  $0$
47. Let  $w = \frac{1-iz}{z-i}$ . If  $|w|=1$ , which of the following must be true?  
 (A)  $z$  lies inside the unit circle  
 (B)  $z$  lies on real axis  
 (C)  $z$  lies on imaginary axis  
 (D)  $z$  lies outside the unit circle  
 (E)  $Re z < 0$
48. For  $|z| \geq 2$ , if  $\left|z + \frac{1}{z}\right| \geq k$ , the minimum possible value of  $k$  is  
 (A)  $1/2$  (B)  $3/2$  (C)  $2$   
 (D)  $5/2$  (E)  $3$
49. Let  $\cot \theta = -5/12$  where  $\frac{\pi}{2} < \theta < \pi$ . Then the value of  $\sin \theta$  is  
 (A)  $-\frac{12}{13}$  (B)  $-\frac{5}{13}$  (C)  $\frac{12}{13}$   
 (D)  $\frac{5}{13}$  (E)  $\frac{7}{13}$
50. The value of  $\tan \frac{\pi}{8}$  is  
 (A)  $\sqrt{2}$  (B)  $-\sqrt{2}$  (C)  $\sqrt{2}-1$   
 (D)  $1-\sqrt{2}$  (E)  $-1-\sqrt{2}$

---

Space for rough work

51. In an A.P., if 5<sup>th</sup> term is  $\frac{1}{7}$  and 7<sup>th</sup> term is  $\frac{1}{5}$ , then the sum of first 35 terms is  
(A) 9 (B) 18 (C) 36  
(D) 72 (E) 83
52. In a G.P.,  $1, \frac{1}{2}, \frac{1}{4}, \dots$ , when the first  $n$  number of terms are added, the sum is  $\frac{1023}{512}$ . Then the value of  $n$  is  
(A) 10 (B) 12 (C) 14  
(D) 16 (E) 18
53. If A.M. and G.M. of the roots of a quadratic equation are 8 and 5 respectively, then the quadratic equation is  
(A)  $x^2 + 8x + 5 = 0$  (B)  $x^2 - 16x + 10 = 0$  (C)  $x^2 - 16x + 25 = 0$   
(D)  $x^2 + 8x + 25 = 0$  (E)  $x^2 + 10x + 15 = 0$
54. Given that the equation  $x^2 - (2a + b)x + \left(2a^2 + b^2 - b + \frac{1}{2}\right) = 0$  has two real roots. The value of  $b$  is  
(A) 1 (B) 2 (C) -1  
(D) -2 (E) 0

---

Space for rough work

55. If  ${}^5P_r = {}^6P_{r-1}$ , then the value of  $r$  is  
(A)  $r = 1$  (B)  $r = 5$  (C)  $r = 3$   
(D)  $r = 2$  (E)  $r = 4$
56. If  ${}^nC_{2017} = {}^nC_{2016}$ , then  ${}^nC_{4033}$  equals  
(A) 1 (B) 2016 (C) 2017  
(D) 2033 (E) 2019
57. The image of the point  $P(2,1)$  on the straight line  $2x - 3y + 1 = 0$  is  
(A)  $\left(\frac{1}{13}, \frac{25}{13}\right)$  (B)  $\left(\frac{15}{13}, \frac{25}{13}\right)$  (C)  $\left(\frac{18}{13}, \frac{25}{13}\right)$   
(D)  $\left(\frac{21}{13}, \frac{25}{13}\right)$  (E)  $\left(\frac{11}{13}, \frac{15}{13}\right)$
58. If the centre of the circle inscribed in a square formed by the lines  $x^2 - 8x + 12 = 0$  and  $y^2 - 14y + 45 = 0$  is  $(a, b)$ , then  $a + b$  is  
(A) 11 (B) 9 (C) 7  
(D) 5 (E) 4

---

Space for rough work



59. The equation of the directrix of the parabola  $y^2 + 4y + 4x + 2 = 0$  is
- (A)  $x = -1$  (B)  $x = 1$  (C)  $x = 3/2$   
 (D)  $x = -3/2$  (E)  $x = 2$
60. The foci of the hyperbola  $\frac{x^2}{\cos^2 \alpha} - \frac{y^2}{\sin^2 \alpha} = 1$  are
- (A)  $(\pm 1, 0)$  (B)  $(\pm \alpha, 0)$  (C)  $(0, \pm 1)$   
 (D)  $(0, \pm \alpha)$  (E)  $(1, \pm \alpha)$
61. The domain of definition of the function  $f(x) = \frac{\log_3(x+7)}{x^2 - 5x + 6}$  is
- (A)  $(-7, \infty) \setminus \{3, 2\}$  (B)  $(-3, \infty) \setminus \{3, 2\}$  (C)  $(-7, \infty) \setminus \{3\}$   
 (D)  $(-3, \infty) \setminus \{3\}$  (E)  $(-5, \infty) \setminus \{3\}$
62. Let  $f(x) = 3x - 5$ . The inverse of  $f$  is given by
- (A)  $\frac{1}{3x-5}$  (B)  $\frac{x+5}{3}$  (C)  $\frac{x}{3} - \frac{1}{5}$   
 (D)  $\frac{x}{3} + \frac{1}{5}$  (E)  $\frac{3}{x-5}$

---

Space for rough work

63. Let  $R = \{(a,b) : a \leq b^2\}$  be a relation on the set of all real numbers. Then  $R$  is
- (A) symmetric but not transitive  
 (B) transitive but not symmetric  
 (C) both symmetric and transitive  
 (D) neither symmetric nor transitive  
 (E) having finite range
64. A unit vector  $\vec{b}$  is coplanar with  $i + j + 2k$  and  $i + 2j + k$  and is perpendicular to  $i + j + k$ . Then  $\vec{b} \cdot i$  equals
- (A) 0 (B) 1 (C)  $\frac{3}{2}$   
 (D) 2 (E) 4
65. Suppose  $\alpha i + \alpha j + \gamma k$ ,  $i + k$  and  $\gamma i + \gamma j + \beta k$  are coplanar where  $\alpha$ ,  $\beta$  and  $\gamma$  are positive constants. Then the product  $\alpha \beta$  is
- (A)  $\gamma$  (B)  $\gamma^2$  (C)  $2\gamma$   
 (D)  $2\gamma^2$  (E)  $3\gamma$
66. The area of the triangle whose vertices are  $A(1, -1, 2)$ ,  $B(2, 1, -1)$  and  $C(3, -1, 2)$  is
- (A)  $\sqrt{7}$  (B)  $\sqrt{11}$  (C)  $\sqrt{13}$   
 (D)  $\sqrt{15}$  (E)  $\sqrt{10}$

---

Space for rough work

67. Let  $f(x) + 2f\left(\frac{1}{x}\right) = \frac{1}{x} - 5$ . Then  $\left| \int_1^2 3f(x) dx \right|$  equals
- (A)  $2 + \ln 2$  (B)  $2 - \ln 2$  (C) 2  
 (D)  $3 \ln 2$  (E)  $\ln 2$
68. The value of  $\lim_{n \rightarrow \infty} \left[ \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{6n} \right]$  is
- (A)  $\ln 3$  (B)  $\ln 6$  (C)  $e^3$   
 (D)  $e^6$  (E)  $\ln 2$
69. Let  $f(x)$  be differentiable and  $\int_0^{t^2} x f(x) dx = \frac{1}{2}t^4$  for all  $t$ . Then the value of  $f(17)$  is
- (A) 17 (B) 1 (C)  $1/17$   
 (D)  $17/2$  (E) 19
70. The value of the definite integral  $\int_0^{2\pi} \sqrt{1 + \sin \frac{x}{2}} dx$  is
- (A)  $\frac{1}{4}$  (B)  $\frac{1}{2}$  (C)  $\frac{3}{4}$   
 (D) 1 (E)  $\frac{5}{4}$
71. Let  $f(x) = |x - 2|$  and  $g(x) = f(f(x))$ . Then derivative of  $g$  at the point  $x = 5$  is
- (A) 1 (B) 2 (C) 4  
 (D) 5 (E) 0

---

Space for rough work



72. Let  $f(x) = \sin x - \cos x$ . Then the value of  $\log_{x \rightarrow \infty} \frac{f(x) - f\left(\frac{\pi}{2}\right)}{x - \frac{\pi}{2}}$  is

- (A) 0 (B)  $\frac{1}{2}$  (C)  $\frac{1}{\sqrt{2}}$   
 (D) 1 (E)  $\sqrt{2}$

73. Let  $A = \begin{pmatrix} \alpha & 0 \\ 1 & 1 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & 0 \\ 5 & 1 \end{pmatrix}$  be two matrices where  $\alpha$  is a real number.

Then

- (A)  $A^2 = B$  for some  $\alpha$  (B)  $A^2 \neq B$  for any  $\alpha$  (C)  $A^2 = -B$  for some  $\alpha$   
 (D)  $|A^2| \neq |B|$  for any  $\alpha$  (E)  $A = -B$  for some  $\alpha$

74. The values of  $k$  for which the system

$$(k+1)x + 8y = 0$$

$$kx + (k+3)y = 0$$

has unique solution, are

- (A) 3, 1 (B) -3, 1 (C) 3, -1  
 (D) -3, -1 (E) 1, -1

---

Space for rough work

75. If  $M$  and  $N$  are square matrices of order 3 where  $\det(M) = 2$  and  $\det(N) = 3$ , then  $\det(3MN)$  is  
 (A) 27 (B) 81 (C) 162  
 (D) 324 (E) 121
76. If the lines  $\frac{x+3}{-3} = \frac{y-1}{k} = \frac{z-5}{5}$  and  $\frac{x+1}{-1} = \frac{y-2}{2} = \frac{z-5}{5}$  are coplanar, then the value of  $k$  is  
 (A) 1 (B) 2 (C) 3  
 (D) 4 (E) 5
77. A plane passes through the point  $P(1, -2, 1)$  and is perpendicular to two planes  $2x - 2y + z = 0$  and  $x - y + 2z = 4$ . Then the equation of the plane is  
 (A)  $x + y + 1 = 0$  (B)  $x - y + 1 = 0$  (C)  $x + 2y + 1 = 0$   
 (D)  $x - 2y + 1 = 0$  (E)  $x - y - 1 = 0$
78. The differential equation which represents the family of curves  $y^2 = 2c(x + \sqrt{c})$  where  $c > 0$ , is of  
 (A) order 2 (B) degree 2 (C) order 3  
 (D) degree 3 (E) degree 1
79. The number of solutions of the differential equation  $\frac{dy}{dx} = y^{1/3}$  which are passing through the origin, is  
 (A) 0 (B) 1 (C) 2  
 (D) 3 (E) 5

---

Space for rough work

80. If  $\frac{dy}{dx} = \frac{2}{x+y}$  and  $y(1) = 0$ , then  $x + y + 2$  equals
- (A)  $3e^{\left(\frac{y}{2}\right)}$  (B)  $2e^{\left(\frac{y}{2}\right)}$  (C)  $e^{\left(\frac{y}{2}\right)}$   
 (D) 0 (E)  $5e^{\left(\frac{y}{2}\right)}$
81. The length of the latus rectum of the parabola  $(x+2)^2 = -14(y-5)$  is
- (A) 7 (B) 14 (C) 21  
 (D) 28 (E) 17
82. One of the foci of the hyperbola  $\frac{x^2}{9} - \frac{y^2}{16} = 1$  is
- (A) (3, 0) (B) (4, 0) (C) (5, 0)  
 (D) (9, 0) (E) (2, 0)
83. If the circles  $x^2 + y^2 - 8x - 6y + c = 0$  and  $x^2 + y^2 - 2y + d = 0$  cut orthogonally, then  $c + d$  equals
- (A) 6 (B) 4 (C) 2  
 (D) 0 (E) 1
84. The points with position vector  $60\hat{i} + 3\hat{j}$ ,  $40\hat{i} - 8\hat{j}$  and  $a\hat{i} - 52\hat{j}$  are collinear if
- (A)  $a = -10$  (B)  $a = 40$  (C)  $a = 20$   
 (D)  $a = 10$  (E)  $a = -40$

---

Space for rough work



85. The area enclosed within the curve  $|x| + |y| = 1$  is

- (A) 1 (B)  $\sqrt{2}$  (C)  $\frac{3}{2}$   
(D)  $2\sqrt{2}$  (E) 2

86. The unit vector in the direction of the vector  $\overline{AB}$  if  $A=(-2, -1, 3)$  and  $B=(1, 1, 0)$  is  $\alpha i + \beta j + \gamma k$ , then  $\alpha + \beta$  is

- (A)  $\frac{3}{\sqrt{22}}$  (B)  $\frac{5}{\sqrt{22}}$  (C)  $\frac{-3}{\sqrt{22}}$   
(D)  $\frac{-5}{\sqrt{22}}$  (E)  $\frac{2}{\sqrt{22}}$

87. If  $\begin{pmatrix} 3x-y & x+3y \\ 2x-z & 2y+z \end{pmatrix} = \begin{pmatrix} 7 & 9 \\ 5 & 5 \end{pmatrix}$ , then  $x+y+z$  equals

- (A) 3 (B) 6 (C) 9  
(D) 12 (E) 11

88. If the product  $abc=1$ , then the value of the determinant

$$\begin{vmatrix} -a^2 & ab & ac \\ ba & -b^2 & bc \\ ac & bc & -c^2 \end{vmatrix} \text{ is}$$

- (A) 1 (B) 2 (C) 3  
(D) 4 (E) 5

---

Space for rough work

89. If  $(x, y, z)$  is the solution of the equations

$$4x + y = 7$$

$$3y + 4z = 5$$

$$5x + 3z = 2$$

Then the value of  $x + y + z$  equals

(A) 8

(B) 6

(C) 3

(D) 0

(E) 1

90. If  $\begin{pmatrix} e & f \\ g & h \end{pmatrix}$  is the inverse of the matrix  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$  where  $ad - bc = 1$ , then  $g$  equals

(A)  $c$

(B)  $-c$

(C)  $b$

(D)  $-b$

(E)  $d$

91. If  $f: \mathbb{R} \rightarrow \mathbb{R}$  is a function defined by  $f(x) = x^2$ , then which of the following is true?

(A)  $f$  is 1-1 but not onto

(B)  $f$  is onto but not 1-1

(C)  $f$  is neither 1-1 nor onto

(D)  $f$  is both 1-1 and onto

(E)  $f^{-1}: \mathbb{R} \rightarrow \mathbb{R}$  exists

---

Space for rough work

92. Consider the set  $A = \{1, 2, 3\}$  along with the relation  $R = \{(1, 1), (2, 2), (1, 2), (2, 1), (3, 3)\}$ . Which of the following statements is **true**?
- (A) The relation is symmetric but not transitive  
 (B) The relation is transitive but not symmetric  
 (C) The relation is neither symmetric nor transitive  
 (D) The relation is both symmetric and transitive  
 (E) The relation is a function
93. If  $(-\sqrt{3} - i)^{30} = -4^k$ , then the value of  $k$  is
- (A) 15 (B) 20 (C) 25  
 (D) 30 (E) 60
94. If  $\omega$  is an imaginary cube root of unity, then  $(1 + \omega - \omega^2)^7$  is equal to
- (A)  $128\omega$  (B)  $-128\omega$  (C)  $128\omega^2$   
 (D)  $-128\omega^3$  (E)  $-128\omega^2$
95. The value of  $\left[\cos\frac{\pi}{8} + i\sin\frac{\pi}{8}\right]^4$  is
- (A)  $-i\pi$  (B)  $i\pi$  (C)  $i$   
 (D)  $-i$  (E)  $\pi$
96. If  $\arg(\bar{z}_1) = \arg(z_2)$ , then
- (A)  $z_2 = kz_1^{-1}, (k > 0)$  (B)  $z_2 = kz_1, (k > 0)$  (C)  $|z_2| = |\bar{z}_1|$   
 (D)  $z_1 = z_2$  (E)  $|z_2| = |z_1|$

---

Space for rough work



97. The value of  $\tan \left[ \sin^{-1} \frac{5}{13} + \cot^{-1} \frac{4}{3} \right]$  is

(A)  $26/11$

(B)  $56/33$

(C)  $63/41$

(D)  $65/43$

(E)  $32/13$

98. If  $\tan^{-1} x + 2 \cot^{-1} x = \frac{\pi}{3}$ , then the value of  $x$  is

(A)  $-\sqrt{3}$

(B)  $-\sqrt{2}$

(C)  $\sqrt{2}$

(D)  $\sqrt{3}$

(E)  $\sqrt{5}$

99. Which of the following is not a solution of the equation

$$3 \tan^2 \theta - \sin \theta = 0?$$

(A)  $n\pi$

(B)  $n \frac{\pi}{2}$

(C)  $n\pi + (-1)^n \frac{\pi}{6}$

(D)  $0$

(E)  $\pi$

---

Space for rough work

100. If  $\sqrt{\frac{y}{x}} + \sqrt{\frac{x}{y}} = 1$ , then  $\frac{dy}{dx}$  equals
- (A)  $\sqrt{\frac{y}{x}}$  (B)  $\sqrt{\frac{x}{y}}$  (C)  $\frac{y}{x}$   
(D)  $\frac{x}{y}$  (E)  $xy$
101. If  $x = \frac{3t}{1+t^3}$  and  $y = \frac{3t^2}{1+t^3}$ , then  $\frac{dy}{dx}$  at  $t = 1$  equals
- (A)  $-6$  (B)  $-1$  (C)  $1$   
(D)  $6$  (E)  $4$
102. The equation of the normal to the curve given by  $x^2 + 2x - 3y + 3 = 0$  at the point  $(1, 2)$  is
- (A)  $3x + 4y - 11 = 0$  (B)  $3x - 4y + 11 = 0$  (C)  $-3x + 4y - 11 = 0$   
(D)  $3x - 4y - 11 = 0$  (E)  $-3x - 4y - 11 = 0$
103. A point of inflection of the curve given by  $y = x^3 - 6x^2 + 12x + 50$  occurs when
- (A)  $x = 2/3$  (B)  $x = 3/2$  (C)  $x = 2$   
(D)  $x = 3$  (E)  $x = 0$

---

Space for rough work

104. The value of the integral  $\int_0^{\frac{\pi}{2}} \log \tan \theta \, d\theta$  is

- (A) 0 (B) 1 (C)  $\frac{\pi}{2}$   
(D)  $\log 2$  (E) 2

105. The area enclosed between the curve  $y = 11x - 24 - x^2$  and the line  $y = x$  is

- (A)  $\frac{1}{3}$  (B)  $\frac{3}{4}$  (C) 1  
(D)  $\frac{4}{3}$  (E)  $\frac{1}{2}$

106. The solution of the differential equation  $\frac{dy}{dx} = \frac{y^2}{x}$  passing through the point (1, -1) is

- (A)  $\frac{1}{y} + \log x = 0$  (B)  $\frac{1}{y} - \log x = 0$  (C)  $y + \log x = 0$   
(D)  $y - \log x = 0$  (E)  $y \log x = 0$

107. The maxima and minima of the function  $2x^3 - 15x^2 + 36x + 10$  occur respectively at

- (A)  $x = 1, x = 3$  (B)  $x = 2, x = 1$  (C)  $x = 3, x = 2$   
(D)  $x = 1, x = 2$  (E)  $x = 2, x = 3$

---

Space for rough work



108. In a class of 100 students, there are 70 boys whose average marks in a subject are 75. If the average marks of the complete class is 72, then what is the average of the girls?
- (A) 73 (B) 85 (C) 68  
(D) 74 (E) 65
109. Let  $x_1, x_2, \dots, x_n$  be  $n$  observations such that  $\sum x_i^2 = 400$  and  $\sum x_i = 80$ . Then a possible value of  $n$  is
- (A) 15 (B) 10 (C) 9  
(D) 12 (E) 18
110. If  $M$  and  $N$  are events such that  $P(M \cup N) = \frac{3}{4}$ ,  $P(M \cap N) = \frac{1}{4}$ ,  $P(\bar{M}) = \frac{2}{3}$ , then  $P(\bar{M} \cap N)$  is
- (A)  $\frac{15}{12}$  (B)  $\frac{3}{8}$  (C)  $\frac{5}{8}$   
(D)  $\frac{1}{4}$  (E)  $\frac{5}{12}$

---

Space for rough work

111. Cards marked with numbers 2 to 105 are placed in a box and mixed. One card is chosen at random. The probability that the number on the card is less than 15 is  
(A)  $\frac{1}{8}$  (B)  $\frac{1}{9}$  (C)  $\frac{7}{8}$   
(D)  $\frac{8}{9}$  (E)  $\frac{2}{7}$
112. An urn contains 4 black, 5 white and 6 red balls. One ball is drawn at random. The probability that it is not black is  
(A)  $\frac{4}{15}$  (B)  $\frac{9}{15}$  (C)  $\frac{11}{15}$   
(D)  $\frac{13}{15}$  (E)  $\frac{14}{15}$
113. In a chess tournament, assume that your probability of winning a game is 0.3 against level 1 players, 0.4 against level 2 players and 0.5 against level 3 players. It is further assumed that among the players 50% are at level 1, 25% are at level 2 and the remaining are at level 3. The probability of winning a game against a randomly chosen player is  
(A) 0.275 (B) 0.375 (C) 0.225  
(D) 0.325 (E) 0.125
114. A man repays a loan of Rs. 3250 by paying Rs. 20 in the first month and then increases the payment by Rs.15 every month. The number of months it takes to clear the loan is  
(A) 20 (B) 25 (C) 35  
(D) 40 (E) 10

---

Space for rough work

115. The coefficient of  $x^3$  in the expansion of  $\left(x^2 - \frac{2}{x}\right)^6$  is

- (A) -160 (B) -80 (C) -40  
(D) 0 (E) -10

116. If the equation of the sphere through the circle

$$x^2 + y^2 + z^2 = 5; 2x + 3y + 4z = 5 \text{ and through the origin is}$$

$$x^2 + y^2 + z^2 - 2x - 3y - 4z + C = 0 \text{ then the value of } C \text{ is}$$

- (A) 1 (B) -1 (C) 0  
(D) 5 (E) 2

117. The equation of the plane containing the lines

$$\frac{x+1}{3} = \frac{y+3}{5} = \frac{z+5}{7} \text{ and } \frac{x-2}{1} = \frac{y-4}{3} = \frac{z-6}{5} \text{ is}$$

- (A)  $x + 2y + z = 0$  (B)  $x - 2y + z = 0$  (C)  $x - 2y - z = 0$   
(D)  $x + 2y - z = 0$  (E)  $2y - x - z = 0$

---

Space for rough work



118. A value of  $c$  for which the conclusion of mean value theorem holds for the function  $f(x) = \log_e x$  on the interval  $[1, 3]$  is

- (A)  $8\log_3 e$                       (B)  $\frac{1}{2}\log_e 3$                       (C)  $\log_3 e$   
(D)  $\log_e 3$                       (E)  $2\log_3 e$

119. From 4 men and 6 ladies a committee of five is to be selected. The number of ways in which the committee can be formed so that men are in majority is

- (A) 68                      (B) 156                      (C) 60  
(D) 72                      (E) 66

120. The degree of the differential equation  $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}} = l \frac{d^2y}{dx^2}$  is

- (A) 1                      (B) 2                      (C) 3  
(D) 4                      (E) 5

---

Space for rough work



**PAPER – II MATHEMATICS VERSION-B1**

1	C	31	A	61	A	91	C
2	B	32	B	62	B	92	D
3	B	33	B	63	D	93	A
4	E	34	B	64	A	94	E
5	E	35	C	65	B	95	C
6	C	36	C	66	E	96	A
7	B	37	B	67	A	97	B
8	B	38	E	68	B	98	A
9	A	39	E	69	B	99	B
10	A	40	E	70	A	100	C
11	C	41	C	71	A	101	B
12	A	42	C	72	D	102	A
13	D	43	B	73	B	103	C
14	D	44	B	74	D	104	A
15	B	45	C	75	C	105	D
16	D	46	D	76	A	106	E
17	E	47	B	77	A	107	E
18	A	48	B	78	D	108	E
19	A	49	C	79	D	109	E
20	A	50	C	80	A	110	E
21	B	51	B	81	B	111	A
22	E	52	A	82	C	112	C
23	A	53	C	83	A	113	B
24	A	54	A	84	E	114	A
25	D	55	E	85	E	115	A
26	B	56	A	86	B	116	C
27	E	57	C	87	B	117	B
28	E	58	A	88	D	118	E
29	E	59	C	89	C	119	E
30	D	60	A	90	B	120	B

*Handwritten signature*  
2/3/15



collegebatch.com

click to campus

## KEAM 2019 Question Paper with Solution

Kerala Engineering Architecture Medical Entrance Exam

KEAM 2019 Question Paper with Solution - Physics & Chemistry	Page No. 2 to 34
KEAM 2019 Question Paper with Solution - Mathematics	Page No. 35 to 67

Download more KEAM Previous Year Question Papers: [Click Here](#)